

AD _____

Award Number: DAMD17-00-1-0317

TITLE: BRCA2 and Genome Integrity

PRINCIPAL INVESTIGATOR: Li-Kuo Su, Ph.D.

CONTRACTING ORGANIZATION: The University of Texas M.D. Anderson
Cancer Center
Houston, Texas 77030

REPORT DATE: July 2001

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20011128 220

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 2001	3. REPORT TYPE AND DATES COVERED Annual (1 Jul 00 - 30 Jun 01)	
4. TITLE AND SUBTITLE BRCA2 and Genome Integrity			5. FUNDING NUMBERS DAMD17-00-1-0317	
6. AUTHOR(S) Li-Kuo Su, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Texas M.D. Anderson Cancer Center Houston, Texas 77030 E-Mail: lsu@mail.mdanderson.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) <p>The <i>BRCA2</i> tumor suppressor gene has been suggested to play an important role in DNA repair and maintaining genome integrity. Most evidences supporting this hypothesis, however, were obtained from studying mouse embryonic stem cells or embryonic fibroblast. The importance of <i>BRCA2</i> in maintaining genome integrity in human cells is not very clear. We have completed the Task 1, generation of Capan-1 derivatives that conditionally express wild type <i>BRCA2</i>. Capan-1 is the only human cell lines known to not express wild type <i>BRCA2</i>. We have obtained two Capan-1 derivatives that express exogenous wild type <i>BRCA2</i> under the regulation of tetracycline. We have also obtained several Capan-1 derivatives that express exogenous mutant <i>BRCA2</i>, either constitutively or regulated by tetracycline. We have also carried out the first part of the Task 3, characterization of Capan-1 derivatives to genotoxic agents. We examined the sensitivity of wild type <i>BRCA2</i>-expressing Capan-1 derivatives to ionizing radiation and DNA damaging chemicals. Our preliminary results showed that there was no detectable difference in the sensitivity to these treatments between when these cells expressed or did not express the wild type <i>BRCA2</i>.</p>				
14. SUBJECT TERMS BRCA2, DNA repair			15. NUMBER OF PAGES 10	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusions.....	7
References.....	8
Appendices.....	10

Introduction

People carrying germline mutations of the breast cancer susceptibility gene *BRCA2* have increased risk for breast, ovarian, pancreatic and other types of cancer (Wooster et al., 1994; Wooster et al., 1995; Tavtigian et al., 1996). Mouse cells lacking a functional *Brca2* gene are deficient in repairing DNA damage (Sharan et al., 1997; Connor et al., 1997; Patel et al., 1998; Davies et al., 2001; Moynahan et al., 2001). Capan-1, a human pancreatic cancer cell line, is the only human cell line known to not express wild-type BRCA2. Capan-1 cells carry only a mutant *BRCA2* (6174delT) and expresses a truncated BRCA2 protein (Goggins et al., 1996; Teng et al., 1996; Chen et al., 1998; Su et al., 1998). The *BRCA2* 6174delT mutation is one that found frequently in Ashkenazi Jews and one that clearly predisposes its carriers to a variety of cancers (Berman et al., 1996; Neuhausen et al., 1996; Oddoux et al., 1996; Roa et al., 1996; Tonin et al., 1996; Abeliovich et al., 1997). Capan-1 cells have been shown to be more sensitive to DNA damaging agents than other human cell lines were (Chen et al., 1998; Abbott et al., 1998). However, Capan-1 cells have many additional genetic alternations compared to these other human cell lines, whether the increased sensitivity of Capan-1 cells to genotoxic agents is caused by the lack of functional BRCA2 is not clear. The goals of this study are to investigate whether alternation of the expression of wild-type BRCA2 in human cell lines would alter the ability of these cells to repair their DNA damage. We have accomplished the Task 1 of this project, to establish Capan-1 derivatives that express wild-type BRCA2. We are carrying out the first part of Task 3 of this project, to investigate whether expression of wild-type BRCA2 alters the sensitivity of Capan-1 to DNA damaging agents.

Body

1. Generation of wild-type BRCA2-expressing Capan-1 derivatives

We established tetracycline regulated wild-type BRCA2-expressing Capan-1 derivatives by transfecting a vector expressing tetracycline regulated activator, tTA-IRES-Neo (Yu et al., 1999), together with a vector expressing BRCA2 cDNA under the control of tetracycline regulated promoter. We identify wild-type BRCA2-expressing Capan-1 derivatives by using immunoblotting using the antibody N61, a BRCA2 monoclonal antibody we generated (Su et al., 1998). After screening about 140 clones obtained from two separate transfection experiments, we

isolated two clones, one from each transfection, that expressed wild-type BRCA2 tightly regulated by tetracycline (figure 1).




Figure 1. Expression of wild-type BRCA2 in Capan-1 derivatives.

Two clones of wild-type BRCA2-expressing Capan-1 derivative have been obtained (Clone A and Clone B). Lysates were prepared from cells grown at the uninduced condition (U), of tetracycline-containing media, or at the induced condition, tetracycline-free media, for 1 (I-1) or 2 (I-2) days. BRCA2 proteins were detected by using immunoblotting using the BRCA2 monoclonal antibody N61 (Su et al., 1998). The arrow and the asterisk indicate the wild-type BRCA2 and the Capan-1 endogenous mutant BRCA2 respectively. Lysates prepared from HCT116 and MCF7 cell lines, which expressed wild-type BRCA2, were used as positive control for wild-type BRCA2.

2. Characterization of wild-type BRCA2-expressing Capan-1 derivatives

We investigated whether expression of wild-type BRCA2 altered the sensitivity of Capan-1 cells to γ -radiation. We compared the sensitivity of these Capan-1 derivatives to γ -radiation between expressing and not expressing wild-type BRCA2. Cells were plated in ten cell culture dishes in the presence of tetracycline, therefore the wild-type BRCA2 was not expressed. The wild-type BRCA2 was induced to express in five dishes of these cells two days after plating. Another two days later, cells were irradiated with different doses of γ -radiation. Cells were grown for another six (for clone B) or seven (for clone A) days and the number of surviving cells in each dish were determined. We have done this experiment three times for clone A and two times for clone B, we are performing the third time for clone B. The results show that expression of wild-type BRCA2 did not result in detectable difference in the γ -radiation sensitivity in these Capan-1 derivatives (figure 2).






Figure 2. Sensitivity of wild-type BRCA2-expressing Capan-1 derivatives to γ -radiation.

(A) The outline of the experiment. Five hundred thousand cells were plated in each of ten 60 mm cell culture dish on day 0. Cells were fed with fresh media on day 2, five dishes of cells were fed with tetracycline-free media to induce the expression of wild-type BRCA2. On day 4 cells were fed with fresh respective media in the morning and were irradiated with different doses of γ -radiation in the afternoon. Cells were then fed with fresh respective media every two days. Surviving cells were harvested on day 10 (clone B) or day 11 (clone A) and the number of surviving cells in each dish was determined. (B, C) Surviving curve of Capan-1 derivatives treated with γ -radiation. The results of three separate experiment of clone A (B) and two separate experiment of clone B (C) are shown. U and I indicate cells grown in uninduced and induced conditions, respectively.

Key Research Accomplishments

- Establishment of Capan-1 derivatives that express wild-type BRCA2 under the regulation of tetracycline

Reportable Outcomes

- Development of Capan-1 derivatives that express wild-type BRCA2 under the regulation of tetracycline
- A manuscript describing the effect of expressing wild-type BRCA2 on the growth of Capan-1 cells is being prepared
- An abstract describing the characterizing the sensitivity of these Capan-1 derivatives to genotoxic agents has been submitted to 51st Annual Meeting (year 2001) of American Society of Human Genetics

Conclusions

We have accomplished Task 1 of this project, generation of wild-type BRCA2-expressing Capan-1 derivatives. We are investigating the sensitivity of these Capan-1 derivatives to DNA damaging agents, the first part of Task 3. We are also working on the Task 2, generation of MCF-12A and MCF7 derivatives that do not express wild-type BRCA2. We will continue working on Task 2 and Task 3 in the next year.

References

- Abbott, D. W., Freeman, M. L., and Holt, J. T. (1998). Double-strand break repair deficiency and radiation sensitivity in BRCA2 mutant cancer cells. *J. Natl. Cancer Inst.* 90, 978-985.
- Abeliovich, D., Kaduri, L., Lerer, I., Weinberg, N., Amir, G., Sagi, M., Zlotogora, J., Heching, N., and Peretz, T. (1997). The founder mutations 185delAG and 5382insC in BRCA1 and 6174delT in BRCA2 appear in 60% of ovarian cancer and 30% of early-onset breast cancer patients among Ashkenazi women. *Am. J. Hum. Genet.* 60, 505-514.
- Berman, D. B., Costalas, J., Schultz, D. C., Grana, G., Daly, M., and Godwin, A. K. (1996). A common mutation in BRCA2 that predisposes to a variety of cancers is found in both Jewish Ashkenazi and non-Jewish individuals. *Cancer Res.* 56, 3409-3414.
- Chen, P.-L., Chen, C.-F., Chen, Y., Xiao, J., Sharp, Z. D., and Lee, W.-H. (1998). The BRC repeats in BRCA2 are critical for RAD51 binding and resistance to methyl methanesulfonate treatment. *Proc. Natl. Acad. Sci. U S A* 95, 5287-5292.
- Connor, F., Bertwistle, D., Mee, P. J., Ross, G. M., Swift, S., Grigorieva, E., Tybulewicz, V. L., and Ashworth, A. (1997). Tumorigenesis and a DNA repair defect in mice with a truncating Brca2 mutation. *Nature Genet.* 17, 423-430.
- Davies, A. A., Masson, J. Y., McIlwraith, M. J., Stasiak, A. Z., Stasiak, A., Venkitaraman, A. R., and West, S. C. (2001). Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. *Mol. Cell* 7, 273-282.
- Goggins, M., Schutte, M., Lu, J., Moskaluk, C. A., Weinstein, C. L., Petersen, G. M., Yeo, C. J., Jackson, C. E., Lynch, H. T., Hruban, R. H., and Kern, S. E. (1996). Germline BRCA2 gene mutations in patients with apparently sporadic pancreatic carcinomas. *Cancer Res.* 56, 5360-5364.
- Moynahan, M. E., Pierce, A. J., and Jasin, M. (2001). BRCA2 is required for homology-directed repair of chromosomal breaks. *Mol. Cell* 7, 263-272.
- Neuhausen, S., Gilewski, T., Norton, L., Tran, T., McGuire, P., Swensen, J., Hampel, H., Borgen, P., Brown, K., Skolnick, M., Shattuck-Eidens, D., Jhanwar, S., Goldgar, D., and Offit, K. (1996). Recurrent BRCA2 6174delT mutations in Ashkenazi Jewish women affected by breast cancer. *Nature Genet.* 13, 126-128.
- Oddoux, C., Struwing, J. P., Clayton, C. M., Neuhausen, S., Brody, L. C., Kaback, M., Haas, B., Norton, L., Borgen, P., Jhanwar, S., Goldgar, D., Ostrer, H., and Offit, K. (1996). The carrier frequency of the BRCA2 6174delT mutation among Ashkenazi Jewish individuals is approximately 1%. *Nature Genet.* 14, 188-190.

- Patel, K. J., Vu, V. P., Lee, H., Corcoran, A., Thistlethwaite, F. C., Evans, M. J., Colledge, W. H., Friedman, L. S., Ponder, B. A., and Venkitaraman, A. R. (1998). Involvement of Brca2 in DNA repair. *Mol. Cell* 1, 347-357.
- Roa, B. B., Boyd, A. A., Volcik, K., and Richards, C. S. (1996). Ashkenazi Jewish population frequencies for common mutations in BRCA1 and BRCA2. *Nature Genet.* 14, 185-187.
- Sharan, S. K., Morimatsu, M., Albrecht, U., Lim, D. S., Regel, E., Dinh, C., Sands, A., Eichele, G., Hasty, P., and Bradley, A. (1997). Embryonic lethality and radiation hypersensitivity mediated by Rad51 in mice lacking Brca2. *Nature* 386, 804-810.
- Su, L.-K., Wang, S.-C., Qi, Y., Luo, W., Hung, M.-C., and Lin, S.-H. (1998). Characterization of BRCA2: temperature sensitivity of detection and cell-cycle regulated expression. *Oncogene* 17, 2377-2381.
- Tavtigian, S. V., Simard, J., Rommens, J., Couch, F., Shattuck-Eidens, D., Neuhausen, S., Merajver, S., Thorlacius, S., Offit, K., Stoppa-Lyonnet, D., Belanger, C., Bell, R., Berry, S., Bogden, R., Chen, Q., Davis, T., Dumont, M., Frye, C., Hattier, T., Jammulapati, S., Janecki, T., Jiang, P., Kehrer, R., Leblanc, J.-F., Mitchell, J. T., McArthur-Morrison, J., Nguyen, K., Peng, Y., Samson, C., Schroeder, M., Snyder, S. C., Steele, L., Stringfellow, M., Stroup, C., Swedlund, B., Swensen, J., Teng, D., Thomas, A., Tran, T., Tran, T., Tranchant, M., Weaver-Feldhaus, J., Wong, A. K. C., Shiuya, H., Eyfjord, J. E., Cannon-Albright, L. A., Labrie, F., Skolnick, M. H., Wever, B., Kamb, A., and Goldgar, D. E. (1996). The complete BRCA2 gene and mutations in chromosome 13q-linked kindreds. *Nature Genet.* 12, 333-337.
- Teng, D. H., Bogden, R., Mitchell, J., Baumgard, M., Bell, R., Berry, S., Davis, T., Ha, P. C., Kehrer, R., Jammulapati, S., Chen, Q., Offit, K., Skolnick, M. H., Tavtigian, S. V., Jhanwar, S., Swedlund, B., Wong, A. K., and Kamb, A. (1996). Low incidence of BRCA2 mutations in breast carcinoma and other cancers. *Nature Genet.* 13, 241-244.
- Tonin, P., Weber, B., Offit, K., Couch, F., Rebbeck, T. R., Neuhausen, S., Godwin, A. K., Daly, M., Wagner-Costalos, J., Berman, D., Grana, G., Fox, E., Kane, M. F., Kolodner, R. D., Krainer, M., Haber, D. A., Struwing, J. P., Warner, E., Rosen, B., Lerman, C., Peshkin, B., Norton, L., Serova, O., Foulkes, W. D., Lynch, H. T., Lenoir, G. M., Narod, S. A., and Garber, J. E. (1996). Frequency of recurrent BRCA1 and BRCA2 mutations in Ashkenazi Jewish breast cancer families. *Nature Med.* 2, 1179-1183.
- Wooster, R., Bignell, G., Lancaster, J., Swift, S., Seal, S., Mangion, J., Collins, N., Gregory, S., Gumbs, C., Micklem, G., Barfoot, R., Hamoudi, R., Patel, S., Rice, C., Biggs, P., Hashim, Y., Smith, A., Connor, F., Arason, A., Gudmundsson, J., Ficenec, D., Kelsell, D., Ford, D., Tonin, P., Bishop, D. T., Spurr, N. K., Ponder, B. A. J., Eeles, R., Peto, J., Devilee, P., Cornelisse, C., Lynch, H., Narod, S., Lenoir, G., Egilsson, V., Barkadottir, R. B., Easton, D. F., Bantley, D. R., Futreal, P. A., Ashworth, A., and Stratton, M. R. (1995). Identification of the breast cancer susceptibility gene BRCA2. *Nature* 378, 789-792.

Wooster, R., Neuhausen, S. L., Mangion, J., Quirk, Y., Ford, D., Collins, N., Nguyen, K., Seal, S., Tran, T., Averill, D., Fields, P., Marshall, G., Narod, S., Lenoir, G. M., Lynch, H., Feunteun, J., Devilee, P., Cornelisse, C. J., Menko, F. H., Daly, P. A., Ormiston, W., McManus, R., Pye, C., Lewis, C. M., Cannon-Albright, L. A., Peto, J., Ponder, B. A. J., Skolnick, M. H., Easton, D. F., Goldgar, D. E., and Stratton, M. R. (1994). Localization of a breast cancer susceptibility gene, BRCA2, to chromosome 13q12-13. *Science* 265, 2088-2090.

Yu, J., Zhang, L., Hwang, P. M., Rago, C., Kinzler, K. W., and Vogelstein, B. (1999). Identification and classification of p53-regulated genes. *Proc. Natl. Acad. Sci. USA* 96, 14517-14522.

Appendix:

None.